



PermafrostNet
NSERC | CRSNG



Hosted by



Natural Sciences and Engineering
Research Council of Canada

Conseil de recherches en sciences
naturelles et en génie du Canada

Canada



Carleton
UNIVERSITY

Theme 1: Permafrost characterization and ground ice potential

PermafrostNet 2023
AGM

Victoria, BC

November 19, 2023

Theme 1: Characterization of permafrost.


Objective:

*To improve the understanding of ground-ice loss and its consequences through better **characterization of permafrost** in the **field** and in **laboratories** so that **prediction** can better represent **processes during thaw** and have relevant subsurface input such as **ground-ice content**.*




Specific objectives:

1. Develop and implement a system for handling permafrost data *that can support prediction, evaluation of prediction and analysis of permafrost change* (PINGO)– the database and field data
2. Evaluate and apply methods to predict and measure thermal, hydrologic geochemical and geomechanical behaviour of frozen soil during thaw to support improved simulation
3. Develop a framework for the spatial and stratigraphic syntheses of geotechnical and geological data to *support ground ice map products*




Olefeldt et al. 2017






Completed



Completed



Completed



Co-investigators



Daniel Fortier (Montreal)
co-lead Theme 1
PDF1, PhD6



Duane Froese (Alberta)
co-lead Theme 1
PDF2, PhD1, PhD5, MSc2



Jocelyn Hayley (Calgary)
co-lead Theme 1
PhD3, MSc1



Pascale Roy-Léveillé
(Laval)
PhD4



Toni Lewkowicz
(Ottawa)
PDF3



Stephan Gruber
PhD2



Partners



Steve Kokelj
NTGS
Yellowknife



Stephen Wolfe,
GSC Ottawa



Sharon Smith
GSC Ottawa




Fabrice Calmels
Yukon University



Ashley Rudy
NTGS
Yellowknife



Brendan O'Neill
GSC Ottawa




Peter Morse
GSC Ottawa



Chris
Stevens
SRK/Independent

Progress

1. PINGO Database structure, fields draft complete
 - *Michel Paquette, Samuel Gagnon, Nick Brown*
2. Ground Ice Potential Database
 - Databases– Yukon, **Mackenzie Valley PIN (Omid's talk, Alex's poster)**, Nunavik
 - Presently ~50,000 ice measurements from ~ 13,000 boreholes primarily Mackenzie Valley
3. Regional studies
 - Hudson Bay Lowlands– **Tabitha Rahman**
 - Mackenzie valley corridor- **Alexandre Chiasson**
 - Mackenzie Mountains and subarctic hillslopes and landslides- **Joe Young**
 - Polar Desert- **Withdrew**



O'Neill et al. 2018

Progress– Next practices

4. Permafrost characterization


- Non-destructive and digital archives
 - Computed tomography- *Mahya Roustaei and Joel Pumple*
 - Multi-sensor core logging- *Joel Pumple, Mahya Roustaei PACS Lab*
- Dielectric methods -- *Hosein Fereydooni Started 2022*
- Geomechanical properties– *Khatereh Roghangar and Zakieh Mohammadi*

5. Electrical resistivity *Teddi Herring*



Mahya Roustaei and Joel Pumple Theme 1 CT and MSCL

Non-destructive methods – GEOTEK (visible ice and ρ)




Multiple rapid non-destructive results

A High resolution core images

B Bulk density (^{137}Cs gamma source)

C Magnetics




Mahya Roustaei and Joel Pumple Theme 1 CT and MSCL

Non-destructive methods – Industrial Ct Scanner- (VIC, EIC, ρ , ρ_s)

Nikon XT H 225 ST



Geotek MSCL





2 papers 2022 CGS Meetings, 1 paper in press The Cryosphere, 2 papers in review

Ground ice potential– progress

1. Databases– Yukon, Mackenzie Valley, ITH, Nunavik
 - Abundance of GWC. (75,000+)
2. PACS Lab Permafrost Index Properties – defines relations between gravimetric water content (GWC) and Excess Ice (**Omid's talk tomorrow**)

Omid Ansghari, Mahya Roustaei, PACS lab Research Associates




Spectral Induced Polarization Field measurements

Hosein Fereydooni

Measurement Sites


- Haines Junction – Pingo
- Carmacks – Buried ice
- Takhini – Retrogressive Thaw Slump
- ❖ Two orthogonal 2-metre-profiles

Equipment



➤ FUCHS III+

A Unit Control, A Current Unit, two Potential Units, A Reference Unit



A Study of Effective Parameters in Thermal Modeling Impacting Permafrost Response to Climate Warming

30 Pages • Posted: 17 Aug 2023

[Khatereh Roghangar](#)

affiliation not provided to SSRN

[Jocelyn L. Hayley](#)

affiliation not provided to SSRN

Abstract

Abstract : Climate warming is causing significant changes in the Arctic, leading to increased temperatures and permafrost instability. The harmonic active layer has been shown to be affected by climate change, where warmer ground surface temperatures result in progressive permafrost thaw and a deepening active layer. This study assessed the effects of critical parameters on permafrost ground response to climate warming using the fifth phase of the Coupled Model Intercomparison Project (CMIP5) and TEMP/W software. We analyzed variations in depth, climate scenarios, water content, and soil types to predict the depth of the active layer and settlement in the future using the soil characteristics along Hudson Bay Railway corridor. The results indicate

Analyzed the effects of thermal modelling parameters (i.e., water content, soil type, model depth, climate scenario and permafrost coverage) on climate-driven permafrost thaw and settlement using the CMIP5 and TEMP/W software

A Framework for Predicting Thaw Settlement in Permafrost Regions

Zakieh Mohammadi

Regional Scale Assessment

Status:

Completed and published.



Local Scale Assessment

Creating predictive tools/methods for thaw settlement by accounting for the unique characteristics of fine-grained, coarse-grained, and highly organic soils.


Status:

- Integrated thaw consolidation data into a unified database (data is ready to be published).
- Work on coarse-grained and organic soils completed (Papers submitted)
- Work on fine-grained sediments in progress.

Creating the Framework

Integrating diverse components to establish a framework for assessing thaw settlement across various scales (Next step)


Past forest fires condition permafrost slopes for failure in the central Mackenzie Valley, NWT



Recent increases in permafrost landslide frequency (278%) and magnitude (602%) between 2004 and 2020 reveal a permafrost landscape in geomorphic transition


More than 80% of permafrost landslides occur in areas burned in the 1990s, indicating the long-term effect of fire on permafrost slope stability

Compounding effects of legacy thermal disturbance from fires and climate drivers of thaw (increasing air temperature and precipitation) are likely to increase slope instability of warm permafrost



Young, J. M., Alvarez, A., van der Sluijs, J., Kokelj, S. V., Rudy, A., McPhee, A., et al. (2022). Recent intensification (2004–2020) of permafrost mass-wasting in the central Mackenzie Valley foothills is a legacy of past forest fire disturbances. *Geophysical Research Letters*, 49, e2022GL100559.

Variability in Permafrost Environment in the Central Mackenzie Valley




Two papers are currently in preparation.

- 1) Chiasson, A., Alvarez, A., van der Sluijs, J., Andersen, B., Kokelj, S., Rudy, A., La Farge-England, C. & Froese, D.G. **Morphology, dynamics and setting of dendritically-drained peat plateaus of the central Mackenzie Valley, Northwest Territories.**
- 2) Chiasson, A., Alvarez, A., van der Sluijs, J., Kokelj, S., Rudy, A., & Froese, D.G. **Investigation of Dendritically-Drained Peat Plateaus: An Analysis of Permafrost Peatland Degradation Over the Past 70 Years in the central Mackenzie Valley, Northwest Territories.**

1. The research has provided new insights into the formation and evolution of peatlands in the central Mackenzie valley, allowing us to better understand the factor that contribute to the variability of peatlands morphology.
2. Peatlands are one of the largest carbon sinks on Earth, and understanding the dynamics of dendritic peat plateaus can help us better predict and mitigate the effects of climate change.
3. The preservation and restoration of dendritic peat plateaus, which are covered in Reindeer Lichens, could not only have a positive impact on carbon sequestration and climate change mitigation efforts, but also provide an important source of nutrition for Caribou and other wildlife.

Tabatha Rahman (Laval) PhD Hudson Bay Lowlands

- In progress:
 - Writing 1st paper on ice-wedge morphology, volume, and distribution in the Barrens of the Hudson Bay Lowlands, northern Manitoba
 - Water isotope analyses on ground-ice samples at PACS lab (Duane's lab)
 - Planning winter fieldwork
- Completed since September:
 - Calculated preliminary ice-wedge volume for the Barrens
 - Collected 297 permafrost samples from 49 boreholes, equivalent to 111.26 m of cores
 - AMS ^{14}C -dated basal peat at 5 study sites
 - CT-scanned all permafrost cores



CPERS Database Update (Teddi Herring, Theme 1)

Created a database of ERT surveys of permafrost

Data can be searched and visualized using the

web interface

The screenshot shows the PermafrostNet web interface. On the left is a sidebar with navigation links: ABOUT CPERS, CPERS COLLECTIVE, MAP OF SURVEYS, LITERATURE REVIEW, DATA PROCESSING, CONTRIBUTE DATA, DATA POLICY, and CONTACT US. The main content area is titled "ERT Surveys of Permafrost" and includes a note about full plotting functionality. Below this are filter sections for "Contains data in date range" (09/22/2008 to 07/05/2022), "Location", "Landform", "Disturbance", and "PI". A map of Canada shows survey locations with colored dots. A detailed information box for "Beaver River Burn" is displayed, including site details, survey dates, and metadata. A 2D ERT plot titled "Beaver River Burn 2: 2018-07-18" shows resistivity data with a color scale from 30 to 404 ρ (mS) and a depth of 10 m.


Data are archived long-term in

Nordicana D

The screenshot shows the Nordicana D database website. The header includes a navigation bar with links: Résumé / Abstract, Carte Map, Collaborateurs Contributors, Remerciements Acknowledgements, Versions, Sites, Documentation, and Téléchargement Download. The main content area is titled "La base de données canadienne des relevés de résistivité électrique du pergélisol" (The Canadian Permafrost Electrical Resistivity Survey Database (CPERS)). It includes a section for "CPERS Collective" and a "Résumé / Abstract" section. The abstract describes the CPERS database as a collection of electrical resistivity tomography (ERT) datasets collected in permafrost environments. It also includes a "Citation des données / Data citation" section with the following text: "CPERS Collective 2023. La base de données canadienne des relevés de résistivité électrique du pergélisol, v. 1.0 (2010-2022). Nordicana D121, doi: 10.5885/45855XD-DC0883ABD6094288." Below the text is a map of Canada showing survey locations with colored dots.

Partnerships

- Many local community connections with field programs (Sahtu, ISR, HBL)
- NWT Geological Survey– Thermokarst Collective Mapping Synthesis papers
 - Kokelj and Rudy led papers forthcoming engaged several network investigators and graduate students-
- Geological Survey of Canada
 - Wolfe and O'Neill engaged with network investigators and graduate students– Canadian thermokarst database and MS for the last 16,000 years



Training and Progress

- Student projects proceeding well and training aspects are being met or exceeded
- Ground Ice Potential database—
~50,000 measurements but should move to 75,000 by February or so
 - Omid Ansghari's presentation and Alex Chiasson's Poster

